


587.

44















Melin Court Waterfall, Vale of Neath. From a Photograph by C. H. Waring.

The Ferns of Wales, by Edmond Young.

LIBRARY  
NEW YORK  
BOTANICAL  
GARDEN.








THE  
FERNS OF WALES.

BY  
EDWARD YOUNG.

LIBRARY  
NEW YORK  
BOTANICAL  
GARDEN.



Reath:

PRINTED AND PUBLISHED BY THOMAS THOMAS.

1856.



fQK89  
.Y68



## Introduction.

LIBRARY  
NEW YORK  
BOTANICAL  
GARDEN.

IT is generally considered that there are forty species of ferns found in the British Isles; of these thirty-four are met with in Wales. Of the remaining six, one (*Gymnogramma leptophylla*) is confined to the Channel Islands; two are found only in Scotland, (*Cystopteris montana* and *Pseudathyrium alpestre*); two (*Aspidium cristatum* and *Aspidium rigidum*) are found in only two or three counties in the North of England; and the remaining one (*Trichomanes speciosum*) only in Ireland.

From this it will be seen, that although the present Work illustrates the ferns of Wales in particular, yet that it embraces nearly every species found in Great Britain,—and all of those which are the most beautiful, and therefore most likely to be cultivated in the fernery.

Although the present Work gives what are generally called thirty-four species, yet several of these so nearly resemble one another as to render it difficult to decide whether they are not merely varieties: these will be pointed out in their places, and it will be seen that the differences are so minute, as to make them scarcely worthy of being elevated from varieties into species.

The tendency of cultivators has lately been to multiply varieties, and to give them names as if they were separate species, thus extending the nomenclature of Ferns in an unnecessary and complex manner.

In the rich fern districts of Wales the varieties are infinite; but in the present Work the specimens given are, as nearly as possible, the types of the several species; the varieties being briefly noticed in the descriptions.

The directions for the cultivation of ferns, and the formation and position of the fernery, are given from the best authorities, and from the author's own experience.

WAINCYRCH, NEATH,  
OCTOBER, 1855.

JAN 1 - 1907



## Directions for the Cultivation of Ferns.

THE principal points to be attended to, are:—

1. *Situation.* A North aspect, tolerably well shaded from the direct rays of the sun, and not subject to the drip of trees.

2. *Soil.* For most ferns the soil should be a compost of peat, loam, decayed leaf mould, and sand. Some require old mortar to be mixed with the compost, and some prefer a clayey soil; these will be especially noticed in the descriptions. In all cases, small pieces of brick, sandstone, or charcoal, about the size of hazel-nuts, should be mixed with the soil to keep it open.

3. *Moisture.* All ferns like moisture; therefore, great care should be taken that they be constantly supplied.

4. *Drainage.* This is one of the most essential points; for although ferns require so much moisture, yet they are materially injured if it be allowed to stagnate; therefore great care should be taken to effect a complete drainage. In pots the bottoms should be filled up at least two inches with pieces of broken bricks, pots, or stones, and above these should be spread a layer of Sphagnum, (a moss always to be found in boggy places, and which never gets mouldy), to prevent the soil washing or settling down among the drainage.

5. *Removing.* Great care should be taken in removing ferns not to injure the roots, but to retain as much as possible of the native soil with the plant. The best time is during the months of March and April.

6. *Growing from Seed.* A very interesting method of propagating from the seed or spores is recommended by Mr. Moore, in his "Handbook of British Ferns:"—"Half fill some shallow wide-mouthed pots with broken crocks, and on this put a layer of about two inches of turfy peat soil and mellow loam, mixed with soft sand-stone broken in small lumps of the size of peas; this compost should not be much consolidated. Next, shake or brush very gently over a sheet of white paper a frond of the species to be propagated; the fine brown dust thus liberated consists of the spores, in greater or less quantity, intermixed more or less with the spore cases. This dust is to be regularly and thinly scattered over the rough surface of the soil, which is immediately to be covered with a bell-glass, large enough to fit down closely within the pot. The pots are at once to be set in feeders and these are to be filled up with water; they may either be placed under a hand-glass in a cold frame, or in a greenhouse or stove, as may be most proper. The first indications of germination will consist in the appearance of little semi-transparent green scales. The supply of water must be kept up, and the glasses should be tilted on one side for a short time every day, and ultimately entirely removed,—the pots still being retained under a hand-glass. After a week or two they may be taken up, carefully separated, and potted singly in small pots. The young plants should be still kept under a hand-glass until established, and then gradually inured to the degree of exposure proper for the mature plants."

The following plan for growing ferns can be strongly recommended, having been found by the author to answer well:—

His fernery consists of a small house twelve feet by five, (facing the North); the roof is of glass, painted white, to prevent the entrance of too much light; nearly the whole of the front is occupied by a door, which during the summer consists of canvas stretched on a wooden frame; and in the winter of board, to keep the frost out; the inside is filled up with a combination of limestone, sandstone, and old trunks of trees, in which the ferns are planted; on the top of this rock-work there is a pipe perforated with very small holes, and connected with a tank of water: by this means a constant drip is obtained, and particular care has been taken to effect a thorough drainage.

This method would answer equally well where heat is used, and would be far preferable to potting the ferns and arranging them on shelves.



Wardian cases are much improved by the introduction of rock work in miniature, and those ferns which are accustomed to stony situations thrive better in consequence.

### The parts of a Fern are :---

The *Roots*,—which are wiry-like fibres thrown out from the rhizome.

The *Rhizome*, or *Caudex*, is a root-like stem, in some ferns creeping either beneath or upon the surface of the soil or rock upon which it grows, and in others erect and tufted.

The *Fronde* springs from the rhizome, and is a term applied to the stem and the leafy part of the fern, but generally refers to the latter part only, the stem being mentioned separately.

Fronds are either—

1. Simple, or undivided ; as *Scolopendrium vulgare*, page 22.
2. Pinnatifid, or more or less deeply cleft, the divisions being termed lobes ; as *Polypodium vulgare*, page 1.
3. Pinnate, or divided into distinct leaf-like divisions, called pinnæ ; as *Asplenium viride* and *trichomanes*, page 17.
4. Bipinnate, the pinnæ themselves being pinnate,—this second series of pinnæ being called pinnules ; as *Polypodium Dryopteris*, page 3.
5. The pinnæ, pinnatifid ; as *Aspidium oreopteris*, page 9.
6. The pinnules, pinnatifid ; as *Asplenium Filix-fœmina*, page 21.
7. The pinnules, pinnate ; as *Pteris aquilina*, page 23.

The *seed* or *spores* of ferns are generally produced on the under surface of the frond, in little *capsules* or spore cases, surrounded by a jointed, elastic ring, which terminates below in a short stalk : the capsule when ripe splits open, and the ring by a series of jerks throws out the seed. The capsules are clustered together, and the clusters are called *sori*.



*Polypodium Vulgare.*

## COMMON POLYPODY.

**A**LTHOUGH this species is very abundant, it is nevertheless, one of the prettiest and most interesting of ferns, for it is found covering those spots where there is the greatest scarcity of vegetation, such as bare rocks and walls; and when an aged oak begins to lose its vigour, this fern establishes itself on its trunk, as if to compensate for its want of verdure in other parts.

The young fronds make their appearance in May and June, issuing from all parts of the rhizome, except the growing point; they arrive at maturity in September, and remain green until the following spring, except in very exposed places.

The rhizome is thick, branching in all directions, and is covered at first with a dense, brown, wiry membrane, which falls off in the winter, leaving a smooth surface.

The stem is green, and nearly as long as the frond.

The fronds vary in length from two to eighteen inches; they are pinnatifid, the margin of the lobes being more or less serrated or notched.

The seed is produced on the under surface of the frond, and is of a bright yellow or orange colour at first, and brown when ripe.

*Habitat.*

Common on moss-covered trunks of trees, on rock, and walls, shady hedge-banks, and stone covered roofs.

*Varieties.*

This species is subject to several variations, the pinnæ being sometimes bifid or divided near the ends, and sometimes more or less deeply cleft. These varieties are scarce in Wales.

*Culture.*

In transplanting this fern, as much as possible of the moss and decayed matter with which the roots are surrounded should be removed with it. It grows freely in pots, if a light porous soil is used, and the rhizomes kept on the surface.

Mr. Newman, in his "History of British Ferns," mentions a very excellent method of cultivating this species in a green-house, viz., "in a wooden basket, made very open, which should be suspended. The rhizomes should be arranged in the bottom of the basket in such a manner that the fronds and the growing points of the rhizomes may pass through the interstices. They should then be covered with a thin layer of Sphagnum, over which spread a mixture of well decayed leaf mould and silver sand; then arrange a second layer of Sphagnum, and then a second layer of rhizomes, on which carefully fasten wooden cross bars, and the basket will be complete. Immerse the whole in soft water, until it is thoroughly saturated, and then suspend it in its final destination. This should be done in April, before any young fronds have appeared: in June and July young fronds will emerge through all the apertures in the basket, and will arrange themselves gracefully around it."







## *Polygona vulgaris.*

### COMMON POLYPOD.

ALTHOUGH this species is very abundant, it is nevertheless one of the most interesting of ferns, for it is found covering those spots where there is the least soil, such as bare rocks and walls; and when an seedling begins to grow, it sends out runners from its trunk, as if to compensate for its want of verdure in other parts. The young plants make their appearance in May and June, bearing from 22 to 24 leaves. Before the growing point, they arrive at maturity in September, and remain green the following year, and in very exposed places.

The stems of this branched fern in all directions, and is covered all over with scales, which fall off in the winter, leaving a smooth surface.

The stem is green, and nearly as long as the frond.

The fronds are from two to eighteen inches, they are pinnate, the rachis being more or less serrated or notched.

The seed is produced on the under surface of the frond, and is of a bright yellow or orange color at first, and brown when ripe.

### *Cultivation.*

Cultivate on moss-covered trunks of trees, on rock, and walls, shallow beds, and in a covered room.

### *Propagation.*

This species is subject to several varieties. The pinnules are sometimes entire, divided, or deeply notched, and sometimes more or less deeply notched. These varieties are named as follows.

### *var. pinnatifida.*

In propagating this fern, as much as possible of the roots and detached leaves with roots, if some are detached should be preserved with it. If grown freely in water, it is a light porous soil used, and the moisture kept on the surface.

Mr. Newman, in the *Illustrations of British Ferns*, gives a very excellent method of cultivating this species in a greenhouse, viz. "In a small basket, made very open, which should be suspended. The drainage should be arranged in the bottom of the basket in such a manner that the fronds and the growing points of the runners may pass through the perforations. They should then be covered with a thin layer of moss, which should be renewed if it will decay, but could not decay more. The moss should be a variety of *Sphagnum*, and then a second layer of drainage, or which would be better, a third layer of moss, and the basket will be complete. It is never the better to use better soil if it is necessary, and then suspend it in the greenhouse. This should be done in April, May, and June, and the young plants will be produced in June and July, and the plants will grow through all the summer in the basket, and will arrange themselves gradually around it."



















[17]

*Polypodium filipense*

## SEEDLING

VERY RARE to the species just described, this elegant fern seems to share the haunts of man, and to exclude itself in the woods or near the falls and waterfalls of our mountains.

The young fronds appear in May, and develop very rapidly, arriving at perfection in July; they are soon cut off by the early frosts on the approach of winter.

The rhizome is erect, very, and creeping; and does not penetrate deep into the soil.

The leaves are green, very little, and much longer than the leafy part of the frond, and is slightly pinnate.

The fronds are very dense, four to five inches in length; they are pinnate, the pinnae being very small.

The seeds are produced on the under surface of the frond, in circular clusters round the edges of the frond, and are of a brown color.

## HABIT.

It grows in deep woods, and near streams, and even under the spray of waterfalls, in the mountainous districts.

## VARIETY.

This species is subject to very little variation.

## CULTURE.

It is easy to cultivate either in pots or in rock-work, requiring to be plentifully supplied with water, and carefully shaded from the rays of the sun, as too much light is prejudicial.

If placed in pots, the rhizome should be covered with soil; great care should be taken to effect a thorough drainage, and the pots should be placed in basins filled with water. The soil should be light and strong.





[ II ]

*Polypodium Phlegopteris.*

BEECH FERN.

**C**ONTRARY to the species just described, this elegant fern seems to shun the haunts of men, and to seclude itself in the woods, or near the rills and waterfalls of our mountainous districts.

The young fronds appear in May, and develop very rapidly, arriving at perfection in July; but are soon cut off by the early frosts on the approach of winter.

The rhizome is tough, wiry, and creeping; and does not penetrate deep into the soil.

The stem is green, very brittle, and much longer than the leafy part of the frond, and is slightly scaly.

The fronds vary from four to twenty inches in length; they are pinnate, the pinnæ being deeply pinnatifid.

The seed is produced on the under surface of the frond, in circular clusters round the edges of the lobes, and is of a brown colour.

*Habitat.*

Common in damp woods, and near streams, and even under the spray of waterfalls, in the mountainous districts.

*Varieties.*

This species is subject to very little variation.

*Culture.*

It is easy to cultivate either in pots or on rock-work, requiring to be plentifully supplied with water, and carefully shaded from the rays of the sun, as too much light is prejudicial.

If planted in pots, the rhizomes should be covered but lightly with soil: great care should be taken to effect a thorough drainage, and the pots should be placed in feeders filled with water. The soil should be light and strong.



## [ III ]

*Polypodium Dryopteris.*

## O A K F E R N .

**T**HE brilliant green colour, and the delicacy of this little fern, cannot fail to charm the beholder. Like the last species, it seeks seclusion, but in dryer situations.

The young fronds appear in April, each at first resembling three little balls on wires; they are soon developed, and bear seed as early as June, but disappear before winter.

The rhizome is very slender and creeping, and when long established, forms a dense mass.

The stem is long and slender, dark purple and shining, and clothed with a few scales at the base.

The fronds are from four inches to a foot in length, of which the stem forms two thirds; they are divided into three triangular branches, each having a short stem which unite at a point with the main stem; each division of the frond is pinnate, and the pinnæ are pinnate at the base, and pinnatifid at the centre and end.

The seed is produced in clusters on the under margin of the pinnules and lobes, and is of a brown colour.

*Habitat.*

Not so common as the last species, but nevertheless abundant in the mountainous districts. It grows in woods, and in shady, rocky places.

*Varieties.*

This species never seems to vary, unless the next species be considered a variety—which is not at all unlikely, as they resemble one another very closely.

*Culture.*

It is easily cultivated. should have a light soil, be protected from the rays of the sun, and be well supplied with moisture.





## St. John's Wort

1811

**H**ERB. Brit. 1811. p. 100. (1811). This plant is very common in the fields and meadows of the British Isles. It is a small, upright, branched plant, with many small, opposite, lanceolate leaves, and small, yellow, tubular flowers.

The young stems are covered with a soft, downy hair, and the leaves are smooth, or slightly hairy on the underside. The flowers are small, and the fruit is a small, round, black berry.

The leaves are very small, and the flowers are very small. The plant is very common in the fields and meadows of the British Isles. It is a small, upright, branched plant, with many small, opposite, lanceolate leaves, and small, yellow, tubular flowers.

The leaves are very small, and the flowers are very small. The plant is very common in the fields and meadows of the British Isles. It is a small, upright, branched plant, with many small, opposite, lanceolate leaves, and small, yellow, tubular flowers.

The leaves are very small, and the flowers are very small. The plant is very common in the fields and meadows of the British Isles. It is a small, upright, branched plant, with many small, opposite, lanceolate leaves, and small, yellow, tubular flowers.

The leaves are very small, and the flowers are very small. The plant is very common in the fields and meadows of the British Isles. It is a small, upright, branched plant, with many small, opposite, lanceolate leaves, and small, yellow, tubular flowers.

The leaves are very small, and the flowers are very small. The plant is very common in the fields and meadows of the British Isles. It is a small, upright, branched plant, with many small, opposite, lanceolate leaves, and small, yellow, tubular flowers.

The leaves are very small, and the flowers are very small. The plant is very common in the fields and meadows of the British Isles. It is a small, upright, branched plant, with many small, opposite, lanceolate leaves, and small, yellow, tubular flowers.













(17)

*Polypodium Calceolatum*

LINDLEY'S POLYPODIES

The species so nearly resembles the last-mentioned that all the time there can only easily be distinguished for the other.

It usually appears in May, and soon after it is noticed.

The plant is creeping, and smaller than the last species.

The leaves are pale coloured, and much broader than the last, and is clothed with scales.

The fronds are from six to eight or ten inches in length; they are partially three-branched, but the branches are much smaller than the main stem. The colour of the fronds is a dull green, and the lower being covered with very small scales, giving a downy appearance to the whole plant, and the scaly stem, are readily distinguished from the last.

It is produced in clusters on the lower surface of the branches of the last, and is of a

small

is confined to the limestone strata, and is of rare occurrence. It has been found near Brecon, and on the Brecon road; the only other locality recorded is Llanfyllin, in Denbighshire. It is not probable that it grows elsewhere, but has been passed by in mistake for the last species.

Calceol

It is more rare than the last. The soil should be light and enriched with lime or manure. Care should be taken not to give it too much moisture.





[IV]

*Polypodium Calcarenm.*

L I M E S T O N E P O L Y P O D Y .

**T**HIS species so nearly resembles the last-mentioned that at the first glance one may easily be mistaken for the other.

The fronds appear in May, and soon arrive at maturity.

The rhizome is creeping, and stouter than the last species.

The stem is pale coloured, and much longer than the frond, and is clothed with scales.

The fronds are from six to eighteen inches in length; they are partially three-branched, but the lower pinnæ are much smaller than the centre division. The colour of the fronds is a dull green, owing to their being covered with very small stalked glands, giving a downy appearance to the surface; which, and the scaly stem, are excellent distinguishing characters.

The seed is produced in clusters on the under margin of the pinnules and lobes, and is of a brown colour.

*Habitat.*

It is confined to the limestone districts, and is of rare occurrence. It has been found near Merthyr-Tydvil on the Brecon road; the only other locality recorded is Llanferris, in Denbighshire. There is little doubt that it grows elsewhere, but has been passed by in mistake for the last species.

*Culture.*

It will bear more sun than *Dryopteris*. The soil should be light and intermixed with lime or old mortar. Care should be taken not to give it too much moisture.



\* *Woodsia Ilvensis*.

RAY'S WOODSIA.

**T**HIS is the rarest Welsh fern, being found only in one county, and there only in a very few places. It is equally rare as a British species, being found only in two English and three Scotch counties. The young fronds appear in the spring, and die down before winter.

The roots are black and wiry; the rhizome is thick and tufted.

The stem is very distinctly jointed at a short distance from its junction with the rhizome; it is of a yellowish colour when mature, and is covered with scales and hairs, as is also the whole of the frond, especially the under surface, the sori being nearly concealed by them.

The fronds, including the stem, are rarely found more than six inches in length: they are pinnate, and the pinnæ are pinnatifid.

The seed is produced on the under edge of the lobes.

*Habitat.* In the fissures of rocks—in the most bleak and exposed situations. It has been found on Snowdon and Glyder-Vawr, in Caernarvonshire,—but in very small quantities.

*Varieties.* There is a very distinct variety of this species, called *Hyperborea*, which some authors consider a species, but it so much resembles *Ilvensis* in general characteristics, that others have placed it among the varieties. It is equally, if not more, rare than *Ilvensis*, and has been found in the same localities; the fronds are much narrower than those of *Ilvensis*, and the pinnæ are more distant from each other, and are nearly triangular.

*Culture.* The *Woodsia* may be easily cultivated either with or without heat: if without, it should have a free circulation of air. The soil should be light and mixed with small pieces of granite or sandstone; the rhizomes should be wedged between two pieces of stone. The plant should be well supplied with moisture, which should be carefully drained off.

*Ceterach Officinarium*.

SCALE FERN.

**A**s species peculiar for being rarely found in its native habitats, but on old walls and ruins. The young fronds appear in the spring, and remain evergreen for several years.

The roots are very short, and are remarkable for their power of penetrating mortar; the rhizome is tufted.

The stem is short, thick, and covered with narrow pointed scales.

The fronds vary in length from four to eight inches: they are deeply pinnatifid, the divisions being alternate.

The whole of the under surface of the frond is covered with dense scales, beneath which are the capsules in oblong clusters. The colour of the scales at first is nearly white, changing to brown when the frond is mature.

*Habitat.* Abundant in the limestone districts of South Wales, but not so common in North Wales.

*Varieties.* This species never seems to vary.

*Culture.* It is not easily cultivated, which probably arises from the roots having been injured in removing. It should be planted in a shady place, and be well supplied with moisture,—which should be given to the roots only and not to the fronds. The soil should have old mortar and small stones intermixed.

\* The specimens given of *Woodsia Ilvensis* are Norwegian, but are precisely similar to those of Wales, of which it would have been impossible to get sufficient for this work.





## *Alnus glutinosa*

Willow

**A**LNUS is the rarest. Willows here, being found near the water, and there being a great number of them. It is usually here as in the other species. They are found along the river and in the marshes. The young trees appear in the spring and are very tender.

The roots are black and very hard. The bark is thin and smooth.

The stem is very straight, yellowish at first, becoming brown as it grows older. It is of a yellowish colour when mature, and is covered with scales and hairs as in the other species. Especially the under surface, the scales being mostly covered by them.

The leaves, including the stem, are rarely found more than six inches in length. The leaves are pinnate.

The seed is produced on the outer edge of the leaf.

It is the source of water in the great tank and exposed situations. It is found in some places near Hyderabad, in the mountains, but in very small quantities.

There is a very distinct variety of this species, called *hyperborea*, which some consider as a species, but it so much resembles *glutinosa* in general characteristics, that it is placed among the varieties. It is reported to be not more than five inches high, but has been found in some localities. The leaves are much smaller than those of *glutinosa*, and the young ones are more from each other and are nearly regular.

**Culture.** The *Alnus glutinosa* may be easily cultivated either with or without fire. If without fire, it should have a free circulation of air. The soil should be good and mixed with small stones or gravel or sandstone. The distance should be varied between two plants of same. The plants should be well supplied with culture, which should be carefully drained off.

## *Alnus glutinosa*

Willow

**A**LNUS glutinosa is the most common. It is found in its native habitat, but on old walls and in the marshes. The young trees are very tender, and are very susceptible for several years.

The roots are very hard, and are very susceptible for the power of penetrating water.

The bark is thin.

The stem is straight, thin, and covered with narrow pinnate scales.

The leaves are very large, from half a foot to one foot. They are deeply pinnate, the pinnules being alternate.

The seed of the *Alnus glutinosa* is produced in small, white, round, smooth, which are very susceptible to damp weather. The seed of the *Alnus glutinosa* is very small, and is very susceptible to damp weather.

**Culture.** *Alnus glutinosa* may be cultivated either with or without fire. It is found in its native habitat, but on old walls and in the marshes.

**Alnus glutinosa.** The young trees are very tender.

**Alnus glutinosa.** It is very susceptible to damp weather. The seed of the *Alnus glutinosa* is very small, and is very susceptible to damp weather. It should be placed in a warm place, and should be carefully drained off. The seed of the *Alnus glutinosa* is very small, and is very susceptible to damp weather.

**Alnus glutinosa.** The young trees are very tender.

**Alnus glutinosa.** It is very susceptible to damp weather. The seed of the *Alnus glutinosa* is very small, and is very susceptible to damp weather. It should be placed in a warm place, and should be carefully drained off. The seed of the *Alnus glutinosa* is very small, and is very susceptible to damp weather.













*Aspidium Encalypta*

## HOLLY-FERN

This fern is well known, for the fronds are nearly as rigid and prickly as holly. It has very few localities, being only found in the Snowdon district, and there only in the highest and most exposed places.

The fronds appear early in the Spring, and remain evergreen for many years.

The fronds are thick, waxy, and very strong; the rhizome is tufted and covered with scales, as in *Aspidium*, which is very short.

The fronds are from three to eight or ten inches in length; they are pinnate, and the pinnae are very narrow, the serratures terminating in sharp spines.

The sori are produced in large circular clusters, on the under surface of the pinnae, in a regular row, half way between the mid-vein and the margin. It is of a dark brown colour when mature.

## Habitat.

It has been found on Moel Hebog, Glyder Fawr, and on rocks above Llyn Idwal, in Gwynedd.

## Remarks.

This species is the first of a group that are subject to a great deal of variation, and take so many intermediate forms,—which are, as it were, connecting links with one another,—that it is difficult to determine which are the species, and which the varieties. The next link to *Encalypta* is *Lobatum*; by many considered a species. It is not so rigid as *Encalypta*, and is found generally near streams in the mountainous districts.

## Culture.

This fern is rarely found in success in out-of-door ferneries. If potted, it should be wedged between two pieces of stone; the soil should be light and sandy; the roots should be well supplied with moisture, which should be carefully drained off.





[ VII ]

*Aspidium Lonchitis.*

HOLLY FERN.

**T**HIS species is well named, for the fronds are nearly as rigid and prickly as holly. It is a very scarce fern, being only found in the Snowdon district, and there only in the loftiest and most inaccessible places.

The young fronds appear early in the Spring, and remain evergreen for many years.

The roots are black, wiry, and very strong; the rhizome is tufted and covered with scales, as is also the stem, which is very short.

The fronds are from three to eighteen inches in length; they are pinnate, and the pinnæ are deeply serrated, the serratures terminating in sharp spines.

The seed is produced in large circular clusters, on the under surface of the pinnæ, in a regular series, half way between the mid-vein and the margin. It is of a dark brown colour when mature.

*Habitat.*

It has been found on Moel Hebog, Glyder Fawr, and on rocks above Llyn Idwell, in Caernarvonshire.

*Varieties.*

This species is the first of a group that are subject to a great deal of variation, and take so many intermediate forms,—which are, as it were, connecting links with one another,—that it is difficult to determine which are the species, and which the varieties. The next link to *Lonchitis* is *Lobatum*: by many considered a species. It is not so rigid as *Lonchitis*, and is found generally near streams in the mountainous districts.

*Culture.*

This fern is rarely found to succeed in out-of-door ferneries. If potted, it should be wedged between two pieces of stone; the soil should be light and sandy; the roots should be well supplied with moisture, which should be carefully drained off.



[ VIII ]

*Aspidium Aculeatum.*

PRICKLY FERN.

**T**HIS seems to be the next link to the variety—*Lobatum*—just mentioned.

The young fronds appear in April, the ends being curved back in a very graceful manner. They arrive at maturity in July, and remain green till the following spring.

The roots are long, strong, and tough; the rhizome is tufted and woody.

The stem is short, and covered with large scales.

The fronds are from one to three feet in length; they are pinnate, the pinnæ being deeply pinnatifid, and the edges of the divisions prickly.

The seed is produced in circular clusters on the under surface of the pinnæ.

*Habitat.*

Common in hedge-rows, and by the sides of streams.

*Culture.*

It is readily cultivated either in or out of doors. The soil should be tolerably rich.





*Asplenium Nidula*

BLACKLY FERN.

**H**ABIT. — This fern is the most common to the country — *Asplenium* — 2nd mentioned.

The young fronds appear in April, the roots being covered back in a very graceful manner. They arrive at maturity in July, and remain green till the following spring.

The roots are long, slender, and rough, the rhizome is short and woody.

The stem is short, and covered with large scales.

The fronds are from one to three feet in length; they are pinnate, the pinnae being deeply divided, and the edges of the divisions prickly.

The seed is produced in circular clusters on the under surface of the pinnae.

*Asplenium*

Common in hedge-rows and by the sides of ditches.

*Asplenium*

It is readily collected either in wet or dry places. The soil is not so rich as that of the *Asplenium*.













*Aspidium Aculeare.*

## SOFT PRICKLY FERN.

This species so much resembles the last mentioned, that it is almost to be mistaken for it, but the differences are as follows.

The roots and rhizome are similar to those of *Aculeatum*.

The stem is longer, and the fronds are more delicate, being cut off by the watery fronds, and the sori are prickle instead of pinnatifid.

## Habitat.

Common in woods and in hedge-rows.

## Cultivation.

It is as easily cultivated as the last species.





[IX]

*Aspidium Angulare.*

SOFT PRICKLY FERN.

**T**HIS species so nearly resembles the last mentioned, that it is difficult to point out the differences.

Its roots and rhizome are similar to those of *Aculeatum*.

The stem is longer, and the fronds are more delicate, being cut off by the wintry frosts; and the pinnæ are pinnate instead of pinnatifid.

*Habitat.*

Common in woods and in hedge-rows.

*Culture.*

It is as easily cultivated as the last species.



[ X ]

*Aspidium Oreopteris.*

MOUNTAIN FERN.

**T**HIS is by no means an abundant British species, although it occurs plentifully in Wales.

The young fronds appear in May, and die down on the approach of winter; they spring in a circle from the rhizome.

The roots are numerous and wiry; the rhizome is tufted and scaly.

The stem is short and covered with scales.

The fronds are from one to four or five feet in length; they are pinnate, the pinnæ are deeply pinnatifid, and are very short at the base, gradually lengthening to the centre, and diminishing to the top. Over the surface of the under side of the pinnæ are scattered small, yellowish, glandular globules, which, when pressed, emit an odour not at all disagreeable.

The seed is produced in circular clusters on the under margin of the lobes, and is of a brown colour.

*Habitat.*

It occurs on damp heaths, hedge-rows, and in woods.

*Varieties.*

This species never seems to vary.

*Culture.*

Some cultivators find this a difficult species to establish. If planted in pots, they should be large, well drained, and placed in feeders of water; the soil should be clay or yellow loam. The plants should be well shaded.





The seed is produced in the ~~ovary~~ on the under surface of the leaf and is of













[57]

*Aspidium Cheilanthes*

## MARSH FERN

A very local species,—only four Welsh localities being known.

The fronds are of two kinds, barren and fertile; the barren appear in May, and the fertile in June, and both die down on the approach of winter.

The roots are black and fibrous; the rhizome is creeping, nearly smooth and of a dark colour. The stem is very long and slender, but thicker in the fertile fronds.

The barren fronds are from one to two feet in length, the fertile often attain the height of four feet, and are pinnate.

The fertile fronds are much less abundant than the barren ones.

The seed is produced in circular clusters on the under margin of the leaves.

## Habitat.

It is found only in bogs and marshy places. The localities are Llaneris, in Glamorganshire; Llaneris, in Anglesea; Anglesea bog, in Glamorganshire, and Penallt bog, near Treherbert, Glamorganshire.

## Cultivation.

It is subject to no cultivation.

## Remarks.

The species thrives well in cultivation. The soil should be composed of peat and decayed leaf-mould; the rhizome should be placed near the surface of the soil; the much moisture cannot be supplied, and it should be allowed to stagnate.





[ XI ]

*Aspidium Thelypteris.*

MARSH FERN.

**A** very local species,—only four Welsh habitats being known.

The fronds are of two kinds, barren and fertile; the barren appear in May, and the fertile in June, and both die down on the approach of winter.

The roots are black and fibrous; the rhizome is creeping, nearly smooth, and of a dark colour.

The stem is very long and slender, but thicker in the fertile fronds.

The barren fronds are from one to two feet in length, the fertile often attain the height of four feet; both are pinnate.

The fertile fronds are much less abundant than the barren ones.

The seed is produced in circular clusters on the under margin of the lobes.

*Habitat.*

It is found only in bogs and marshy places. The localities are Llanberis, in Caernarvonshire; Llwydeard lake, Pentraeth, in Anglesea; Singleton bog, in Glamorganshire; and Pennalle bog, near Tenby, Pembrokeshire.

*Varieties.*

It is subject to no variation.

*Culture.*

This species thrives well in cultivation. The soil should be composed of peat and decayed leaf-mould; the rhizome should be planted near the surface of the soil; too much moisture cannot be supplied, and it should be allowed to stagnate.



[ XII ]

*Aspidium Filix-mas.*

MALE FERN.

**T**HIS is one of our most abundant species. The young fronds appear early in the Spring, and survive the winter.

The rhizome is tufted and covered with brown scales, as is also the stem, which is short and stout.

The fronds are from one to four feet in length; they are pinnate, the lower pinnæ being pinnate, the upper ones pinnatifid.

The seed is produced in circular clusters, on the under surface of the lobes, half way between the mid-vein and the margin.

*Habitat.*

Abundant everywhere in hedge-rows, and in woods.

*Varieties.*

There is a variety rather abundant in some parts of Wales, in which the pinnæ are pinnate, and the pinnules are deeply serrated: fronds often occur, in which the pinnæ are bifid, or divided near the extremity.

*Culture.*

This species is readily cultivated, requiring a light porous soil.





*Asplenium platyneuron*

Willd. Sp. Pl. 4: 637

*Asplenium platyneuron* is a small fern, growing in wet places. The fronds are dark green, and the leaves are smooth. The rhizome is covered with brown scales, and the stem is erect.

The fronds are bipinnate, and the pinnules are small and narrow. The lower part of the frond is covered with small, dark spots. The seed is small and round, and the root is fibrous.

*Asplenium platyneuron* is a small fern, growing in wet places.

The fronds are bipinnate, and the pinnules are small and narrow. The lower part of the frond is covered with small, dark spots.

The fronds are bipinnate, and the pinnules are small and narrow.















*Aspidium Cuneifolium*

WITHERING'S BROAD LEAF

This is the first of a group of three, which are subject to so much variation, as to be almost allied to each other, that it is difficult to determine which are the species and which the varieties. Of the three so-called species here given, I consider two as the varieties of the other, though I am unable to discover which is the type of the group.

In this species, the young fronds appear in May, and do not appear the winter or autumn fronds. The rhizome is tufted and covered with pale brown, scaly scales, as is also the lower part of the stem, which is stout and rather long.

The fronds are from one to three feet in length; they are bipinnate, the pinnules being deeply serrated, the lobes terminating in sharp spines; the pinnules are opposite, four or five of the lower ones being equal in length, the others narrowing to the extremity.

The seed is produced in circular clusters on the under surface of the pinnules.

*habitat*

Common in woods and in hedgerows.

*varieties*

As before stated, this species is subject to considerable variation.

*Culture*

Best of cultivation in a light, porous soil.





[ XIII ]

*Aspidium Spinulosum.*

WITHERING'S BROAD FERN.

**T**HIS is the first of a group of three, which are subject to so much variation, and are so closely allied to each other, that it is difficult to determine which are the species and which the varieties. Of the three so-called species here given, I consider two to be varieties of the other, though I am unable to discover which is the type of the species.

In this species, the young fronds appear in May, and often survive the winter in sheltered places. The rhizome is tufted and covered with pale brown roundish scales, as is also the lower part of the stem, which is stout and rather long.

The fronds are from one to three feet in length; they are bi-pinnate, the pinnules being deeply notched, the lobes terminating in sharp spines; the pinnæ are opposite, four or five of the lower pairs being equal in length, the others narrowing to the extremity.

The seed is produced in circular clusters on the under surface of the pinnules.

*Habitat.*

Common in woods, and on hedge-rows.

*Varieties.*

As before stated, this species is subject to considerable variation.

*Culture.*

Easy of cultivation in a light, porous soil.



[ XIV ]

*Aspidium Dilatatum.*

B R O A D F E R N .

**N**EXT to our common brake fern, this is the finest species we have.

The fronds appear in May, and often survive the winter.

The rhizome is tufted, and covered with lanceolate scales, the centre being of a dark brown colour, and the edges of a light brown.

Most part of the stem is covered with similar scales; it is stout, and occupies about one-third of the whole length of the fronds, which often measure five feet.

The fronds are bi-pinnate, the lower pinnules being again pinnate, the upper deeply serrated, the lobes terminating in spines. The pinnæ are opposite the first one or two pairs, being considerably shorter than the others, the next very wide, the others gradually narrowing to the extremity.

The seed is produced in circular clusters on the under surface of the pinnules.

*Habitat.*

Common in woods, and on shady banks.

*Varieties.*

Like the last species, this is subject to considerable variation.

*Culture.*

It may be readily cultivated in a light and porous soil.







1893

## Asplenium

Asplenium

The 12 is our common brake fern, this is the first of the ferns we have.

The fronds are in 12, and are very common.

The rhizome is thick, and covered with brownish scales, the leaves being of a dark green color, and the edges of a light brown.

Most part of the rhizome is covered with scales, which are very common, and are of a dark brown color, the whole length of the rhizome, which is very common.

The fronds are bi-pinnate, the lower pinnules being very small, the upper pinnules being very large, and the edges of the fronds being very common.

The seed is produced in similar manner to the other ferns of our genus.

Asplenium

Asplenium

Asplenium

Asplenium

Asplenium

Asplenium















[XV]

*Aspidium Fernaria*

REPERCHED OR BURN'S FERN

*Aspidium* is the smallest of the group just referred to, and the most interesting, owing to its beautiful green colour contrasted with a dark purple stem.

The fronds appear about May, and are cut off by the early frosts.

The rhizome is tufted, and densely covered with long narrow pale brown scales, or is also the case, which is as long as the frond, the scales are jagged at the end; the colour of the stem is dark

The fronds are from one to two feet in length; they are bipinnate, the pinnules being small, and the sub-pinnules deeply pinnatifid; the lower pair of pinnae are much the largest. The shape of the frond is triangular or deltoid, the colour pale green; it has a crisp appearance, caused by the veins being concave, and the edges curled upwards; on this account this species has been named by some authors *Recurvum*; and this crispness renders it particularly distinct as compared to the *hercynicum*.

Spores are produced in circular clusters on the under surface of the fronds.

*Aspidium*

Found in shady, rocky places, but of rare occurrence in South Wales, having been found only at the Glanvort waterfall, in the Vale of Neath. It is by no means abundant in North Wales, being only occasionally in the Snowdon district; but it occurs in the Ffestiog district in a more common manner.

*Aspidium*

There are several varieties mentioned of this species, but none appear to have been found in

*Aspidium*

It grows readily in a heavy soil, requiring plenty of water and shade.





[XV]

*Aspidium Foenisetii.*

HAY-SCENTED OR BREE'S FERN.

**T**HIS is the smallest of the group just referred to, and the most interesting, owing to its beautiful green colour contrasted with a dark purple stem.

The fronds appear about May, and are cut off by the early frosts.

The rhizome is tufted, and densely covered with long narrow pale brown scales, as is also the stem, which is as long as the frond, the scales are jagged at the end; the colour of the stem is dark purple.

The fronds are from one to two feet in length; they are bi-pinnate, the pinnules being again pinnate, and the sub-pinnules deeply pinnatifid; the lower pair of pinnæ are much the largest. The shape of the frond is triangular or deltoid, the colour pale green; it has a crispy appearance, caused by each of the lobes being concave, and the edges curled upwards: on this account this species has been named by some authors, *Recurvum*; and this crispness renders it particularly difficult to prepare for the herbarium.

The seed is produced in circular clusters on the under surface of the pinnules.

*Habitat.*

Found in shady, rocky places, but of rare occurrence in South Wales, having been found only at the Melincourt waterfall, in the Vale of Neath. It is by no means abundant in North Wales, being met with occasionally in the Snowdon district; but it occurs on the Holyhead mountain in some quantity.

*Varieties.*

There are several varieties mentioned of this species, but neither seems to have been found in Wales.

*Culture.*

It grows readily in a loamy soil, requiring plenty of water and shade.



[XVI]

*Cystopteris fragilis.*

BRITTLE FERN.

**T**HIS is one of the prettiest of the small ferns, and remarkable for its delicate and fragile nature.

The fronds appear in spring, and die down towards the end of autumn.

The rhizome is short and creeping. The stem is smooth, about one-third the length of the frond, which varies from six to eighteen inches. The fronds are bi-pinnate, the lower pinnules being deeply pinnatifid, and the upper ones serrated. The seed is produced in circular clusters on the under surface of the pinnules, rather nearer the mid-vein than the margin.

*Habitat.*

Tolerably abundant in mountainous districts in rocky, shady situations.

*Varieties.*

This species is subject to considerable variation, many of the varieties being very interesting, and considered as distinct species by some authors; but I cannot regard them as other than varieties.

*Culture.*

It is readily cultivated either out of doors or in a green-house, requiring moisture and shade.





*Cylopteris fragilis.*

[ABOVE THE FERN]

**T**HIS is one of the prettiest of the small ferns, and remarkable for its delicate nature.

The fronds appear in spring, and die down towards the end of autumn.

The rhizome is short and creeping. The stem is smooth, about one-third of an inch high, which varies from six to eight inches. The fronds are bipinnate, the lower ones deeply pinnatifid, and the upper ones serrated. The seed is produced in numerous small round sori on the underside of the pinnules, rather nearer the mid-vein than the margin.

[FERN]

Tolerably abundant in mountainous districts in rocky, shady places.

[FERN]

This species is subject to considerable variation, many of the varieties being small and much branched, or distinct species, but I cannot say of them more.

[FERN]

It is usually collected at the foot of the mountain, and grows best in the shade.




















[XVII]

\* *Asplenium Septentrionale*.

FORKED SPLEENWORT.

**T**HIS is the first of the interesting genus, *Asplenium* or Spleenwort.  
The fronds appear in April, and remain evergreen.

The roots are strong, and penetrate deeply into the crevices of rocks; the rhizomes are tufted and often matted together in large masses. The fronds are from two to four inches long, and have very much the appearance of grass, but terminate in forks. The seed is produced in elongated clusters on the under surface of the frond, near the forked ends.

*Habitat.*

Very rare. Occurs in the following localities in Caernarvonshire and Denbighshire:—namely, Carnedd Llewellyn, Llyn y Cwm, Pass of Llanberris, Moel Lechog, Bettws y Coed, near Pont y Pair, and at Llan Dethyla.

*Varieties.*

None.

*Culture.*

Being a tender fern, this requires much care in cultivation: it succeeds well in pots in a cold and shaded frame, requiring plentiful moisture, which should be well drained off.

[XVIII]

*Asplenium Ruta-muraria*.

RUE-LEAVED SPLEENWORT.

**T**HIS pretty little fern must be familiar to everybody, as it occurs abundantly on the walls of most old buildings.

The fronds, which are evergreen, appear about May or June.

The roots are black and wiry; the rhizomes are tufted, and are much matted together. The stem is long, and stronger than the frond, the length of the whole being in general three or four inches. The frond is triangular and bi-pinnate. The seed is produced in diamond shaped clusters, which cover the under surface of the pinnules.

*Habitat.*

Common on rocks and old walls.

*Varieties.*

The only variety of any importance is one which many botanists regard as a distinct species, under the name of *A. Germanicum*; it resembles the normal *Ruta-muraria*, but is simply pinnate, instead of being bi-pinnate. It is very rare; and has been found near Llanrwst, and on Moel Lechog, in Caernarvonshire.

*Culture.*

This fern does not thrive if removed from its native walls or rocks; but if it be desired to cultivate, it should be planted amongst broken bricks and old mortar, and be but sparingly supplied with moisture.

\* The specimens of *A. Septentrionale* given in this work were procured by the author in Norway, and are precisely similar to those found in Wales.



[ XIX ]

*Asplenium Trichomanes.*

MAIDENHAIR SPLEENWORT.

**T**HIS is a well-known little fern, and a particular favourite with cultivators.

The fronds make their appearance in April or May, and are persistent through the winter.

The roots are peculiarly adapted for penetrating into almost invisible fissures in rocks. The rhizome is tufted. The stem is short and black, which together with the frond, extends from three to nine inches. The frond is pinnate, the pinnæ being from a quarter to half an inch in length, and oblong. The seed is produced in clusters, which are arranged diagonally to the mid-veins of the pinnæ.

*Habitat.*

On hedge-rows, walls and rocks, everywhere.

*Varieties.*

There is one marked variety, the pinnæ of which are deeply pinnatifid. I have found it near Penrice Castle, in Glamorganshire.

*Culture.*

It is most readily cultivated, requiring no particular directions.



[ XX ]

*Asplenium Viride.*

GREEN SPLEENWORT.

**T**HIS species much resembles the last, but is more interesting from its brilliant green colour and comparative rarity.

The fronds, which are evergreen, appear in May or June.

The roots are delicate and fibrous, the rhizome tufted. The stem is green and about a third as long as the frond, the whole length being from two inches to a foot. The frond is pinnate; the pinnæ are triangular, and notched in the margin. This triangular shape of the pinnæ, and the brilliant green colour of the stem and frond, form the distinctions between this and *A. Trichomanes*. The seed is produced in clusters similar to the last species.

*Habitat.*

Generally in moist shady situations; often under the drip of waterfalls. It occurs in the Snowdon district, on the Brecon Beacon, at Ysgwd Henrhyd Waterfall, and on the Cribbarth Mountain in the Swansea Valley, and under the Upper Cilhepste Waterfall, in the Vale of Neath.

*Varieties.*

I have not met with any.


*Culture.*

This species thrives well in greenhouses, if planted in pots or on rockwork, and well supplied with moisture.



















[XXI]

*Asplenium Marinum.*

SEA SPIDER FERN.

**T**HIS is a truly marine species, being almost always found within reach of the spray of salt water.

The fronds appear early in the summer, and are evergreen.

The roots are tough, and penetrate into rocks, so as to be removed with difficulty. The rhizome is tufted and hairy. The stem is strong, shining black, and about one-third as long as the frond. The frond is from three to twenty inches long, and pinnae the pinnae being oblong, and more or less serrated. The seed is produced in clusters, disposed like those of *A. Trichomanes*.

## HABIT.


On rocks, and in crevices near the coast. I have found it near Margam Abbey, Glamorganshire, at least three miles from the sea,—but there is reason to believe that the sea has receded from its former margin. It has also been found near Pook, Llan-glyn, in Caernarvonshire, at a still greater distance from the sea, but in this case the sea has been shut out by a dike.

## VARIETIES.

There is no marked variety.

This fern generally requires artificial heat, but may be made to thrive in a cold frame. It should be wedged between two stones, even if potted; the soil should be light and stony.





[ XXI ]

*Asplenium Marinum.*

SEA SPLEENWORT.

**T**HIS is a truly marine species, being almost always found within reach of the spray of salt water.

The fronds appear early in the summer, and are evergreen.

The roots are tough, and penetrate into rocks, so as to be removed with difficulty. The rhizome is tufted and hairy. The stem is strong, shining black, and about one-third as long as the frond. The frond is from three to twenty inches long, and pinnate; the pinnæ being oblong, and more or less serrated. The seed is produced in clusters, disposed like those of *A. Trichomanes*.

*Habitat.*

On rocks, and in caves along the coast. I have found it near Margam Abbey, Glamorganshire, at least three miles from the sea,—but there is reason to believe that the sea has receded from its former margin. It has also been found near Pont Aber-glaslyn, in Caernarvonshire, at a still greater distance from the sea; but in this case the sea has been shut out by a dike.

*Varieties.*

There is no distinct variety.

*Culture.*

This fern generally requires artificial heat, but may be made to thrive in a cold frame. It should be wedged between two stones, even if potted; the soil should be light and stony.



[XXII]

*Asplenium Lanceolatum.*

HUDSON'S SPLEENWORT.

**T**HIS, like the last species, is generally found within a short distance of the sea-coast, but is by no means abundant.

The fronds, which are also evergreen, make their appearance in May.

The roots are black and penetrating. The rhizome is tufted. The stem is about one-third the length of the frond, and is covered with hair-like scales. The fronds are from three inches to a foot long, and pinnate, the pinnae being either pinnate or pinnatifid, and the edges of the pinnules serrate. The form of the frond is lanceolate, whence the fern derives its name. The seed is produced in circular clusters near the margin of the lobes.

*Habitat.*

On rocks and in caves, near Llanrwst; on rocks above Tremadoc, and on a rock called Careg-y Ymbil, in Pwllheli harbour, in Caernarvonshire; on the left hand side of the road between Barmouth and Dolgelly, about a mile from the former place; and near Aber-glaslyn in Merionethshire; and on the south coast of Ramsay Island, near St. David's, Pembrokeshire.

*Varieties.*

None.

*Culture.*

It thrives well in a pot, either with or without heat, if protected and thoroughly drained, and prefers a rough peaty soil.





Asplenium Cernuatum

WINDY SPIDERWORM

*Asplenium Cernuatum*, like the last species, is generally found within a short distance of the water.

The fronds, which are also evergreen, make their appearance in May. The roots are thick and jointed. The rhizome is covered. The fronds are dark green, and are divided into several long, narrow, linear-lanceolate, and are very stiff, and are very stiff, and are very stiff. The fronds are very stiff, and are very stiff. The fronds are very stiff, and are very stiff. The fronds are very stiff, and are very stiff.

*Asplenium Cernuatum* is a very common fern, and is found in many places. It is a very common fern, and is found in many places. It is a very common fern, and is found in many places. It is a very common fern, and is found in many places. It is a very common fern, and is found in many places. It is a very common fern, and is found in many places.

*Asplenium Cernuatum* is a very common fern, and is found in many places. It is a very common fern, and is found in many places. It is a very common fern, and is found in many places. It is a very common fern, and is found in many places. It is a very common fern, and is found in many places.













*Reverend Father to the Right*

*My dear Father, I have the honor to acknowledge the receipt of your letter of the 10th inst.*

Yours of the 10th inst. has been duly received. I am sorry to hear that you are not well, and I hope that you will soon be able to return to your duties. I am, Sir, very respectfully,  
Your obedient servant,  
J. H. [Name]

The above is a copy of the letter which I have just received from you.

The letter is dated the 10th inst. and is addressed to me. The letter is in French and is very interesting. It contains many facts which are of great value to me. I am, Sir, very respectfully,  
Your obedient servant,  
J. H. [Name]

Yours

On this [date] I have the honor to acknowledge the receipt of your letter of the 10th inst.

Yours

This letter is very interesting and contains many facts which are of great value to me. I am, Sir, very respectfully,  
Your obedient servant,  
J. H. [Name]

Yours

It is really a pleasure to hear from you and I am sure that you will find much interest in the work.





[XXIII]

*Asplenium Adiantum-nigrum.*

BLACK MAIDENHAIR SPLEENWORT.

**T**HIS species very much resembles the last, but can readily be distinguished by the form, which is triangular; the lower pinnæ being the largest, whilst in *A. Lanceolatum*, the largest pinnæ are near the centre.

The fronds appear in the beginning of the summer, and are evergreen.

The roots are strong and fibrous. The rhizome is tufted. The stem is black and shining, and nearly as long as the frond, the whole measuring from four to eighteen inches. The frond is bi-pinnate, the lower pinnules being pinnatifid, the upper ones serrate. The seed is produced in very dense linear clusters, near the mid-rib, and is nearly black when mature, whence the name of the fern.

*Habitat.*

On walls, rocks, and stony hedge-rows, everywhere.

*Varieties.*

This fern varies much according to the situation, the lobes being in some cases more narrow and acute, in others more broad and obtuse.

*Culture.*

It is readily cultivated, and is a handsome ornament to rockwork.

[ XXIV ]

*Asplenium Filix-femina.*

L A D Y - F E R N .

**W**HILST one of our most abundant, this is also one of our most elegant ferns.

It is one of the first to appear in the spring, but dies down on the approach of winter.

The rhizome is tufted, and often grows to a considerable length. The stem is thick, short, and covered with scales. The frond is from one to three feet in length. It is bi-pinnate, and the pinules are deeply serrated. The seed is produced in clusters near the mid-rib.

*Habitat.*

Abundant in hedge-rows and in woods.

*Varieties.*

This species is subject to great variation, but neither of the varieties seems, according to my observation, sufficiently marked to deserve a distinct name.

*Culture.*

It is readily cultivated either in or out of doors.





*Asplenium Filix-foemina*

LYNCH

THIS is one of our most abundant, this is the one of our most elegant ferns.

It is one of the first to appear in the spring, but dies down on the approach of winter. The rhizome is tufted, and often grows to a considerable length. The stem is erect, covered with scales. The frond is from one to three feet in length. It is bipinnate, the scales are deeply serrated. The seed is produced in numbers near the middle.

*Asplenium*

Abundant in hedgerows and in woods.

*Asplenium*

This species is subject to great variation, but neither of the varieties is so much observed, sufficiently marked to deserve a distinct name.

*Asplenium*

It is readily cultivated in a common garden.















[XXX]

### *Salicaria Unifida*

PLANT. R. 1000.

The appearance of the first or second division from the plant, the first being one undivided leaf.

The young fronds appear in the spring, and remain unopened for many years.

The rhizome is tufted. The stem is prostrate, the length of the frond, black, and covered with bristly scales. The frond is from six to ten inches long, and entirely undivided. The seed is produced in oblong clusters on the under surface, and diagonally to the mid-vein.

Leaves

On shady boggy-places, in wet fields, and occasionally on walls, abundant.

Leaves

There are several very interesting varieties. The fronds are sometimes very much divided at the base, and sometimes crisped at the edges.

Leaves

It is easily cultivated.



[XXV]

*Scolopendrium Vulgare.*

HART'S TONGUE.

THE appearance of this fern is very different from the generality, the frond being one undivided leaf.

The young fronds appear in the spring, and remain evergreen for many years.

The rhizome is tufted. The stem is one-third the length of the frond, black, and covered with bristly scales. The frond is from six inches to two feet long, and entirely undivided. The seed is produced in oblong clusters on the under surface, and diagonally to the mid-vein.

*Habitat.*

On shady hedge-rows, in damp ditches, and occasionally on walls, abundant.

*Varieties.*

There are several very interesting varieties. The fronds are sometimes very much divided at the top, and sometimes crisped at the edges.

*Culture.*

It is easily cultivated.



[ XXVI ]

*Pteris Aquilina.*

COMMON BRAKES.

**T**HIS is the largest as well as commonest of all ferns, and must be familiar to all.

It appears in the spring, and dies down on the approach of winter.

The rhizome is creeping and velvety. The stem is stout and long, and is remarkable for the disposition of the veins, which, when cut across, form a figure like an oak tree. The frond is tripinnate, and triangular in shape, and often attains a height of eight feet. The seed is produced in a continuous line round the margins of the lobes.

*Habitat.*

Common everywhere.

*Varieties.*

This species seems rarely to vary.

*Culture.*

It is so common that it will hardly ever be desirable to transplant it, but should it be required for the fernery, it must be removed with great care during winter, when the rhizomes are dormant.





[XXVI]

*Alnus Aquatica*

[XXVII]

**A**QUATIC is the name of a water plant, which grows in swamps and bogs.

It appears in the spring, and dies down in the approach of winter.

The rhizome is creeping, and white. The stem is three feet long, and a small quantity of the plant which, when cut, grows in a few days an inch long, pinnate, and trifoliate, is sharp, and will stand upright at eight feet. The plant is continuous from the margin of the bog.

[XXVIII]

[XXIX]

[XXX]

[XXXI]

It is a very common plant, and may be seen in swamps and bogs. It is a very useful plant, and is used in many ways. It is a very common plant, and may be seen in swamps and bogs. It is a very useful plant, and is used in many ways.













*Argemone Oleosa*

*Argemone Oleosa*, the Black-cherry Thistle.

*Argemone Oleosa* is the most common of the Argemone species in the British Isles. It appears in the early spring, and is found in the most fertile soils.

The roots are slender, and the leaves are small and deeply lobed. The stems are erect, and the flowers are large and single. The petals are of a deep red color, and the seeds are black and round.

*Roots.*

Occurs on all high mountains in North Wales, and in South Wales, is found only at the base of the mountains.

*Stems.*

*Leaves.*

*Flowers.*

If planted in a garden or in pots, it should be kept in a warm place, and the soil should be light and sandy.

*Properties.*

*Argemone Oleosa*

*Argemone Oleosa*

*Argemone Oleosa* is the most common of the Argemone species in the British Isles, and is generally found in the most fertile soils.

It appears in the early spring, and is found in the most fertile soils.

The roots are slender, and the leaves are small and deeply lobed. The stems are erect, and the flowers are large and single. The petals are of a deep red color, and the seeds are black and round.

*Roots.*

Occurs on all high mountains in North Wales, and in South Wales, is found only at the base of the mountains.

*Stems.*

The plant is subject to but little variation.

*Leaves.*

It should be planted in a well-draining soil, and the soil should be light and sandy.





[ XXVII ]

*Cryptogramma Crispa.*

ROCK BRAKE, OR PARSLEY FERN.

**T**HIS fern seems to take the place of the Common Brakes on mountains. It appears in May or June, and dies before the winter.

The roots are fibrous; the rhizomes are creeping and closely matted together. The stem is long, straw-coloured and smooth. The fronds are of two sorts—fertile and barren. The barren fronds are bi-pinnate, the pinnules being pinnatifid and notched. The fertile fronds are tri-pinnate, and bear the seed in circular clusters between the mid-rib and the margin.

*Habitat.*

Common on all high mountains, in North Wales; but in South Wales, is found only at Aberdare.

*Varieties.*

None.

*Culture.*

If planted on rockwork or in pots, it should be tightly wedged between two stones, and the soil should be light and stony.

[ XXVIII ]

*Adiantum Capillus-Veneris.*

MAIDENHAIR.

**T**HIS is the most beautiful of the British ferns, and is generally prized by cultivators. It appears in May, and remains green through the winter if sheltered.

The roots are wiry; the rhizome short and creeping. The stem is long, shining and black. The frond is a brilliant green, triangular and bi-pinnate; the pinnules being fan-shaped with notched margins. The seed is produced in clusters round the margin, which is reflexed and covers them.

*Habitat.*

Under Dunraven Castle, on Barry Island, and at a few other places on the south-eastern coast of Glamorganshire.

*Varieties.*

This fern is subject to but little variation.

*Culture.*

It should be planted in a soil containing some portion of lime. It requires abundant moisture and should be well protected.

[XXIX]

*Blechnum Boreale.*

HARD FERN.

**T**HIS is rather a striking fern, and by no means rare.

The fronds, which are of two sorts, fertile and barren, spring up in May; the former die before winter, the latter are evergreen.

The roots are long and wiry; the rhizome is tufted. The stems of the barren fronds are very short, those of the fertile are long and stout. The fronds grow in a whorl, the barren ones being outside and recumbent; the fertile ones within them and erect, and much fewer in number. All the fronds are deeply pinnatifid, so as to be almost pinnate. The seed is produced in two lines, one each side of the mid-rib. The fertile fronds have a peculiarly crisp nature, which renders them difficult to prepare for the herbarium.

*Habitat.*

Common in damp hedge-rows, and on the side of hills.

*Varieties.*

The apex is sometimes much divided, and forms the variety *Ramosum* of some authors. I have met with one root bearing barren fronds only, the lobes of which were deeply notched.

*Culture.*

This fern should be cultivated in a clayey soil, and abundantly supplied with water.





Worms Worm

1800 1820

**W**ORMS is a common name for many different species of insects.

The worms are found in all parts of the world, and are very common in the soil.

The worms are very useful to the farmer, as they eat the rotting matter in the soil, and turn it into manure. They also eat the roots of the plants, and do much damage to the crops.

1800

1800 1820 1840 1860 1880 1900

1800

The worms are very common in the soil, and are very useful to the farmer.

1800

The worms are very common in the soil, and are very useful to the farmer.












155





*Hymenophyllum Tunbridgense.*

TUNBRIDGE FILMY FERN.

**T**HIS and the following are the smallest British ferns, and are remarkable for their moss-like appearance.

It seems to have no particular time for springing up, as new and withered fronds may always be found mixed together in every mass.

The roots are wiry and adhere strongly to the surface of damp rocks. The rhizomes are creeping and much matted together. The stem is about half as long as the frond, and the whole rarely attains the length of three inches. The frond consists of veins very much branched and covered with a filmy wing, which is serrate. The seed is contained in cup-like involucre, supported on short stalks, between the stem and the branches, which split open at the top when the seed is ripe to allow the capsules to escape.

*Habitat.*

On damp rocks, generally near streams, in all mountainous districts.

*Varieties.*

None.

*Culture.*

This fern is difficult to cultivate, but I have found the following plan to succeed best:—Fill the bottom of a large earthenware saucer (pierced in the bottom) with pieces of broken brick or sandstone; over this place a layer of Spagnum, and then a thin layer of soil composed of peat and sand in equal proportions, in which the fern must be planted. It must be kept covered with a plate of glass and well supplied with water.

*Hymenophyllum Wilsoni.*

WILSON'S FILMY FERN.

**T**HIS species is rather larger than *H. Tunbridgense*, but very much resembles it, so that they are often mistaken for one another; but a close inspection shows them to be distinct.

The same description applies to it in every particular, except that the wings are rarely more than three times divided; that they are drooping while those of *H. Tunbridgense* are erect; and the involucre is pointed and entire at the top, while that of *H. Tunbridgense* is rounded and serrated.

*Habitat.*

The same as *H. Tunbridgense*, but it is rather rarer.

*Varieties.*

None.

*Culture.*

The same as *H. Tunbridgense*.

[ XXXII ]

*Osmunda Regalis.*

ROYAL OR FLOWERING FERN.

**T**HIS fern is remarkable for bearing its seed in a sort of crown at the top of the frond, whence, probably, it derives its name.

It appears in May and dies before the winter.

The roots are long and penetrate deep into the ground. The rhizome is tufted and very large, sometimes rising considerably out of the ground. The stem is stout, woody, smooth, and about one-third as long as the frond. The frond attains the length of five to seven feet. It is bi-pinnate, the pinnules being generally entire, sometimes serrated. The seed is produced in circular clusters on the changed pinnules of the upper part of the frond. Instances may often be found where a pinnule is only partly changed, the remainder being leafy.

*Habitat.*

Common in swampy places, and on wet hedge-rows.

*Varieties.*

None.

*Culture.*

It should be grown in peat, and be plentifully supplied with water, which should not be drained off.





Common Fern

ROYAL OR FINEST FERN

It is a very common fern for bearing its seed like most of the ferns of the  
country.

It appears in the autumn before the winter.

The fern is a very common fern in the country. The flowers  
are small and white. The fern is a very common fern in the country.  
It is a very common fern for bearing its seed like most of the ferns of the  
country.

Stem

Leaves of the fern are very common in the country.

Stem

Stem

Stem

If the fern is found in the country, it is a very common fern for bearing its seed like most of the ferns of the country.  
It is a very common fern for bearing its seed like most of the ferns of the country.





















*Ophioglossum Vulgatum.*

## ADDER'S TONGUE.

**A**T first sight, this by no means resembles a fern, but is soon distinguished by its seed.

It appears in the spring, and dies in the course of the summer.

The roots are very long and penetrating. The rhizome hardly exists, as the stem almost springs from the roots. The stem is principally underground, and longer than the frond. The frond consists of a single leaf, two or three inches long, and ovate. From within the folds of this leaf, rises a stalk, longer than the leaf, and bearing on its top a spike of a double row of capsules.

*Habitat.*

In swampy fields abundant, in some localities; but I have met with it in only two, namely, at Glyn Clydach and Pen-y-Graig, near Neath, Glamorganshire.

*Varieties.*

A dwarf variety has been described as a separate species, under the name of *O. Lusitanicum*; but I have not heard of its occurrence in Wales.

*Culture.*

As the roots extend to a considerable depth, it is necessary to remove a large sod, in which it must be allowed to remain, supplying it plentifully with water, which should be slowly drained off.

*Botrychium Lunaria.*

## MOONWORT.

**T**HIS is another frequenter of meadow land, and in many respects resembles the last species. It appears about April, and dies down during the summer.

The roots are succulent and brittle, the rhizome a mere point. The stem is mostly underground and about the same length as the frond. The frond is from two to four inches long, and consists of one fertile and one barren leaf. The barren leaf is pinnate, the pinnæ being half-moon shaped. The fertile leaf is bi-pinnate, and covered with globular capsules, which are yellowish brown when ripe.

*Habitat.*

It is found in dryer situations than the last species, but is rather local. I am only acquainted with a few of its localities in the neighbourhood of Neath.

*Varieties.*

A variety is described as *B. Rutaceum*, but I have not heard of its being found in Wales.

*Culture.*

This fern should be cultivated in the same manner as the last.



# Index.


No. of Page.	No. of Fern.		
1 ..	1	Polypodium Vulgare.. .. .	.. Common Polypody.
2 ..	2	Do. Phegopteris. .. .	.. Beech Fern.
3 ..	3	Do. Dryopteris. .. .	.. Oak Fern.
4 ..	4	Do. Calcareum... .. .	.. Limestone Polypody.
5 ..	5	Woodsia Ilvensis. .. .	.. Ray's Woodsia.
" ..	6	Ceterach Officinatum. .. .	.. Scale Fern.
6 ..	7	Aspidium Lonchitis .. .	.. Holly Fern.
7 ..	8	Do. Aculeatum. .. .	.. Prickly Fern.
8 ..	9	Do. Angulare .. .	.. Soft Prickly Fern.
9 ..	10	Do. Oreopteris .. .	.. Mountain Fern.
10 ..	11	Do. Thelypteris. .. .	.. Marsh Fern.
11 ..	12	Do. Filix-mas. .. .	.. Male Fern.
12 ..	13	Do. Spinulosum. .. .	.. Withering's Broad Fern.
13 ..	14	Do. Dilatatum. .. .	.. Broad Fern.
14 ..	15	Do. Fœnisecii .. .	.. Hay-scented, or Bree's Fern.
15 ..	16	Cystopteris Fragilis. .. .	.. Brittle Fern.
16 ..	17	Asplenium Septentrionale. .. .	.. Forked Spleenwort.
" ..	18	Do. Ruta-muraria. .. .	.. Rue-leaved Spleenwort.
17 ..	19	Do. Trichomanes. .. .	.. Maidenhair Do.
" ..	20	Do. Viride .. .	.. Green Do.
18 ..	21	Do. Marinum.. .. .	.. Sea Do.
19 ..	22	Do. Lanceolatum.. .. .	.. Hudson's Do.
20 ..	23	Do. Adiantum-nigrum .. .	.. Black Maidenhair Do.
21 ..	24	Do. Filix-femina .. .	.. Lady-Fern.
22 ..	25	Scelopendrium Vulgare .. .	.. Hart's Tongue.
23 ..	26	Pteris Aquilina. .. .	.. Common Brakes.
24 ..	27	Cryptogramma Crispa .. .	.. Rock Brake or Parsley Fern.
" ..	28	Adiantum Capillus-Veneris .. .	.. Maidenhair.
25 ..	29	Blechnum Boreale .. .	.. Hard Fern.
26 ..	30	Hymenophyllum Tunbridgense .. .	.. Tunbridge Filmy Fern.
" ..	31	Do. Wilsoni .. .	.. Wilson's Do.
27 ..	32	Osmunda Regalis .. .	.. Royal, or Flowering Fern.
28 ..	33	Ophioglossum Vulgatum .. .	.. Adder's Tongue.
" ..	34	Botrychium Lunaria .. .	.. Moonwort.









New York Botanical Garden Library  
QK89 .Y68 f  
Young, Edward/The ferns of Wales gen  
  
3 5185 00089 2966



